

“The effect of vaginal and caesarean delivery on lower urinary tract symptoms: what makes the difference?”: reply to comment by Mahajan et al.

Henriette Jorien van Brummen

Received: 4 March 2007 / Accepted: 4 March 2007 / Published online: 6 April 2007
© International Urogynecology Journal 2007

Dear Editor,

We thank Dr Mahajan and colleagues for their comment on our article [1]. Unfortunately we are not informed whether the cesarean section performed in our study group involved use of a bladder flap. We do recognize the possibility of trauma to the bladder when performing a CS, possibly leading to SUL.

Reference

1. Van Brummen HJ, Bruinse HW, van de Pol G, Heintz APM, van der Vaart CH (2007) The effect of vaginal and caesarean delivery on lower urinary tract symptoms: what makes the difference? Int Urogynecol J 18:133–139 DOI [10.1007/s00192-006-0119-5](https://doi.org/10.1007/s00192-006-0119-5)

This is a reply to the comment that can be found at
<http://dx.doi.org/10.1007/s00192-007-0344-6>.

H. J. van Brummen (✉)
Department of Perinatology and Gynecology,
University Medical Center Utrecht,
Room F05.216, Heidelberglaan 100, P.O. Box 85500,
Utrecht, GA 3508, The Netherlands
e-mail: hbrummen@zonnet.nl